

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/112039/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Vann, Seralynne ORCID: <https://orcid.org/0000-0002-6709-8773> and Nelson, Andrew ORCID: <https://orcid.org/0000-0002-5171-413X> 2018. Anterior thalamic nuclei lesions have a greater impact than mammillothalamic tract lesions on the extended hippocampal system: a reply. *Hippocampus* 28 (10), pp. 691-693. 10.1002/hipo.22953 file

Publishers page: <http://dx.doi.org/10.1002/hipo.22953>
<<http://dx.doi.org/10.1002/hipo.22953>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Anterior thalamic nuclei lesions have a greater impact than mammillothalamic tract lesions on the extended hippocampal system: A reply

Dear Editor,

Perry et al. recently carried out a replication of anterior thalamic nuclei (ATN) and mammillothalamic tract (MTT) lesion studies, looking at the effects of these lesions on watermaze and radial-arm maze tasks. Consistent with previous findings, they found MTT lesions impaired working memory tasks in both the water-maze (Santin et al., 1999; Vann, 2013; Vann and Aggleton, 2003) and radial-arm maze (Nelson and Vann, 2014; Vann, 2013; Vann and Aggleton, 2003). In contrast, rats with MTT lesions were unimpaired on a reference memory task in the water-maze, again consistent with studies showing mammillary body lesions have little or no effect on this task (Santin et al., 1999; Sutherland and Rodriguez, 1989). The authors found more pronounced impairments following ATN lesions, which is again consistent with previous studies that have directly compared ATN and mammillary body lesions (Aggleton et al., 1991; Aggleton et al., 1995; Gaffan et al., 2001) (see also Sutherland and Rodriguez (1989) who tested rats with mammillary body and ATN on the same water-maze task). Those studies found ATN lesions to be more disruptive than mammillary body lesions on standard tasks, including T-maze alternation, but there was greater similarity across the lesion groups when task demands increased or non-spatial cues were removed.


Perry et al.'s behavioural findings are therefore entirely as expected given our current knowledge. However, the authors claim to be testing a model that predicts equivalent impairments following lesions of the MTT and ATN. Furthermore, they erroneously imply that this is a position that we have advanced. For example, in the abstract it is written, **"it is assumed that their (the MTT and ATN) influence on memory is functionally equivalent"**. In the Introduction it is written, **"New experimental evidence suggests a more prominent influence of the MTT and a greater influence of the MB's brainstem connections on the integrity of the extended hippocampal system than was previously recognised (Dillingham et al., 2015; Vann and Nelson, 2015). The latter perspective suggests that equivalent deficits after MTT and ATN injury would be common-place rather than the exception"**; in the Discussion they write, **"recent animal work, however, has shown that brainstem structures influence the memory system upstream via their impact on the MB, which suggests that damage to the MTT afferents to the ATN would produce comparable memory impairments to that found after ATN lesions (Vann, 2013; Dillingham et al., 2015; Vann and Nelson, 2015)."** In setting up the study in this manner the authors have simply created a straw-man argument. This is not a viewpoint we hold and this is not what is written in the papers that are cited; such claims would disregard both anatomical and functional data (e.g., Jankowski et al., 2013). Furthermore, while the tegmental inputs have been shown to be important, it does not follow that MTT and ATN lesion effects would be comparable, i.e., highlighting a role for the tegmental inputs does not automatically reduce the previously identified contributions of other thalamic connections. Given ATN lesions

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1002/hipo.22953

result in retrosplenial/hippocampal deafferentation and the ATN have a number of efferents and afferents that have been shown to support memory, it is self-evident that ATN lesions should produce greater impairments. We would never say otherwise given much of the data to the contrary comes from our own laboratories. In the Dillingham et al. (2015) paper we devote an entire section to the finding that ATN lesions are typically more disruptive than MTT lesions and possible reasons for this dissociation (p.113-114). In the Vann and Nelson (2015) paper we state, “a further implication is that anterior thalamic lesion effects are to some extent driven by the loss of their inputs from mammillary body lesions” (p.170). The ‘to some extent’ makes it clear that we are not espousing that the ATN function can be explained entirely by their mammillary body inputs.

An additional issue is that the authors do not give an accurate representation of the findings from the geometric task in the watermaze. In the Introduction they write, **“For example, MTT lesions in rats were reported to show weak and transient reference memory impairments in the standard water-maze in one report and did not impair geometric learning in a water-maze in another study (Winter et al., 2011; Vann, 2013). By contrast, ATN lesions produce profound deficits on both tasks (Warburton et al., 1999; Wolff et al., 2008; Aggleton et al., 2009; Dumont et al., 2014)”** and again in the Discussion, **“differences between the spatial memory effects of ATN lesions and MTT lesions have been reported using other tasks. Rats with MTT lesions are able to discriminate the fixed location in a geometric learning task at the same rate as controls whereas rats with ATN lesions remain at chance levels (Aggleton et al., 2009; Vann, 2013; Dumont et al., 2014).”** This is misleading. In terms of the geometric task, MTT lesions that *spare the lateral mammillary projections* did not affect task performance (Vann, 2013). In contrast, lateral mammillary body lesions significantly impaired performance on the geometric task (Vann, 2011). Fornix lesions do not affect performance of this geometric task (Aggleton et al., 2009). Therefore, the most parsimonious argument is that this task is particularly sensitive to lesions within the head direction system. So our point was that this is not a distinction between the ATN and MTT/mammillary bodies but a distinction between head direction and non-head direction regions (Aggleton et al., 2009; Vann, 2011; Vann, 2013).

Finally, the authors assessed Zif268 expression in their lesion groups. However, they do not make it clear that their Zif268 results for the MTT lesion group differ from a previous study (Frizzati et al., 2016). The authors make a point that their MTT lesions do not affect Zif268 in deep retrosplenial layers or reduce Zif268 in dysgranular cortex. However, we found decreases in superficial and deep layers in both Rgb and dysgranular retrosplenial cortex (Frizzati et al., 2016). These differences may be due to our more stringent criteria for lesion inclusion or because we [and (Dumont et al., 2012) whose results also differed from those presented by Perry et al.] closely controlled the behavior across our experimental groups. These points aside, there does seem to be a reduction in Zif268 staining in the deep layers of the MTT group when looking at Figure 2b (see areas highlighted in red), although this apparent visual reduction does not correspond with the quantitative analyses, so perhaps this brain section is not representative.



In summary, the authors found a marked spatial impairment on two behavioural tasks following MTT lesions, consistent with previous reports. They found more widespread impairments following ATN lesions, again consistent with previous reports. There is consensus across the field that while the projections from the mammillary bodies to the anterior are important for some aspects of memory, in particular rapid allocentric encoding, other non-mammillary body projections also contribute. It must be the goal of future research to use more nuanced behavioural tasks combined with selective lesions of inputs to the anterior thalamic nuclei to determine the specific roles of these different pathways for memory.

Yours Faithfully,

Seralynne Vann and Andrew Nelson

References

- Aggleton JP, Keith AB, Sahgal A. 1991. Both fornix and anterior thalamic, but not mammillary, lesions disrupt delayed non-matching-to-position memory in rats. *Behav Brain Res* 44:151-61.
- Aggleton JP, Neave N, Nagle S, Hunt PR. 1995. A comparison of the effects of anterior thalamic, mamillary body and fornix lesions on reinforced spatial alternation. *Behav Brain Res* 68:91-101.
- Aggleton JP, Poirier GL, Aggleton HS, Vann SD, Pearce JM. 2009. Lesions of the fornix and anterior thalamic nuclei dissociate different aspects of hippocampal-dependent spatial learning: implications for the neural basis of scene learning. *Behav Neurosci* 123:504-19.

- Dillingham CM, Frizzati A, Nelson AJ, Vann SD. 2015. How do mammillary body inputs contribute to anterior thalamic function? *Neurosci Biobehav Rev* 54:108-19.
- Dumont JR, Amin E, Poirier GL, Albasser MM, Aggleton JP. 2012. Anterior thalamic nuclei lesions in rats disrupt markers of neural plasticity in distal limbic brain regions. *Neuroscience* 224:81-101.
- Frizzati A, Milczarek MM, Sengpiel F, Thomas KL, Dillingham CM, Vann SD. 2016. Comparable reduction in Zif268 levels and cytochrome oxidase activity in the retrosplenial cortex following mammillothalamic tract lesions. *Neuroscience* 330:39-49.
- Gaffan EA, Bannerman DM, Warburton EC, Aggleton JP. 2001. Rats' processing of visual scenes: effects of lesions to fornix, anterior thalamus, mamillary nuclei or the retrohippocampal region. *Behav Brain Res* 121:103-17.
- Jankowski MM, Ronnqvist KC, Tsanov M, Vann SD, Wright NF, Erichsen JT, Aggleton JP, O'Mara SM. 2013. The anterior thalamus provides a subcortical circuit supporting memory and spatial navigation. *Front Syst Neurosci* 7:45.
- Nelson AJ, Vann SD. 2014. Mammillothalamic tract lesions disrupt tests of visuo-spatial memory. *Behav Neurosci* 128:494-503.
- Santin LJ, Rubio S, Begega A, Arias JL. 1999. Effects of mammillary body lesions on spatial reference and working memory tasks. *Behav Brain Res* 102:137-50.
- Sutherland RJ, Rodriguez AJ. 1989. The role of the fornix/fimbria and some related subcortical structures in place learning and memory. *Behav Brain Res* 32:265-77.
- Vann SD. 2011. A role for the head-direction system in geometric learning. *Behav Brain Res* 224:201-6.
- Vann SD. 2013. Dismantling the Papez circuit for memory in rats. *Elife* 2:e00736.
- Vann SD, Aggleton JP. 2003. Evidence of a spatial encoding deficit in rats with lesions of the mammillary bodies or mammillothalamic tract. *J Neurosci* 23:3506-14.
- Vann SD, Nelson AJ. 2015. The mammillary bodies and memory: more than a hippocampal relay. *Prog Brain Res* 219:163-85.